

## A Note on Rational Approximation with Respect to Metrizable Compactifications of the Plane

M. FRAGOULOPOULOU, V. NESTORIDIS

*Department of Mathematics, University of Athens  
Panepistimiopolis, Athens 157 84, Greece  
fragoulop@math.uoa.gr, vnestor@math.uoa.gr*

Presented by Manuel Maestre

Received February 10, 2015

*Abstract:* In the present note we examine possible extensions of Runge, Mergelyan and Arakelian Theorems, when the uniform approximation is meant with respect to the metric  $\varrho$  of a metrizable compactification  $(S, \varrho)$  of the complex plane  $\mathbb{C}$ .

*Key words:* compactification, Arakelian's theorem, Mergelyan's theorem, Runge's theorem, uniform approximation in the complex domain.

AMS *Subject Class.* (2010): 30E10.

### 1. INTRODUCTION

It is well known that the class of uniform limits of polynomials in  $\overline{D} = \{z \in \mathbb{C} : |z| \leq 1\}$  coincides with the disc algebra  $A(D)$ . A function  $f : \overline{D} \rightarrow \mathbb{C}$  belongs to  $A(D)$  if and only if it is continuous on  $\overline{D}$  and holomorphic in the open unit disc  $D$ . It is less known (see [3, 7]) what is the corresponding class when the uniform convergence is not meant with respect to the usual Euclidean metric on  $\mathbb{C}$ , but it is meant with respect to the chordal metric  $\chi$  on  $\mathbb{C} \cup \{\infty\}$ . The class of  $\chi$ -uniform limits of polynomials on  $\overline{D}$  is denoted by  $\tilde{A}(D)$  and contains  $A(D)$ . A function  $f : \overline{D} \rightarrow \mathbb{C} \cup \{\infty\}$  belongs to  $\tilde{A}(D)$  if and only if  $f \equiv \infty$ , or it is continuous on  $\overline{D}$ ,  $f(D) \subset \mathbb{C}$  and  $f|_D$  is holomorphic. The function  $f(z) = \frac{1}{1-z}$ ,  $z \in D$ , belongs to  $\tilde{A}(D)$ , but not to  $A(D)$ ; thus, it cannot be uniformly approximated on  $D$ , by polynomials with respect to the usual Euclidean metric on  $\mathbb{C}$ , but it can be uniformly approximated by polynomials with respect to the chordal metric  $\chi$ .

More generally, if  $K \subset \mathbb{C}$  is a compact set with connected complement, then according to Mergelyan's theorem [10] polynomials are dense in  $A(K)$  with respect to the usual Euclidean metric on  $\mathbb{C}$ . We recall that a function  $f : K \rightarrow \mathbb{C}$  belongs to  $A(K)$  if and only if it is continuous on  $K$  and holomorphic in the interior  $K^\circ$  of  $K$ .

An open problem is to characterize the class  $\tilde{A}(K)$  of  $\chi$ -uniform limits of polynomials on  $K$ .

CONJECTURE. ([1, 6]) Let  $K \subset \mathbb{C}$  be a compact set with connected complement  $K^c$ . A function  $f : K \rightarrow \mathbb{C} \cup \{\infty\}$  belongs to  $\tilde{A}(K)$  if and only if it is continuous on  $K$  and for each component  $V$  of  $K^\circ$ , either  $f(V) \subset \mathbb{C}$  and  $f|_V$  is holomorphic, or  $f|_V \equiv \infty$ .

Extensions of this result have been obtained in [5] when  $K^c$  has a finite number of components and  $K$  is bounded by a finite set of disjoint Jordan curves. In this case, the  $\chi$ -uniform approximation is achieved using rational functions with poles out of  $K$  instead of polynomials. Furthermore, extensions of Runge's theorem are also proved in [5]. Finally a first result has been obtained in [5] concerning an extension of the approximation theorem of Arakelian ([2]).

Instead of considering the one point compactification  $\mathbb{C} \cup \{\infty\}$  of the complex plane  $\mathbb{C}$ , we can consider an arbitrary metrizable compactification  $(S, \varrho)$  of  $\mathbb{C}$  and investigate the analogues of all previous results. This is the content of the present paper.

## 2. PRELIMINARIES

We say that  $(S, \varrho)$  is a *metrizable compactification of the plane*  $\mathbb{C}$ , if  $\varrho$  is a metric on  $S$ ,  $S$  is compact,  $S \supset \mathbb{C}$  and  $\mathbb{C}$  is an open dense subset of  $S$ . Obviously,  $S \setminus \mathbb{C}$  is a closed subset of  $S$ . We say that the points in  $S \setminus \mathbb{C}$  are the points at infinity.

Let  $(S, \varrho)$  be a metrizable compactification of  $\mathbb{C}$  with metric  $\varrho$ . Many such compactifications can be found in [1]. The one point compactification  $\mathbb{C} \cup \{\infty\}$  with the chordal metric  $\chi$  is a distinct one of them. We note that in this case, the continuous function  $\pi : S \rightarrow \mathbb{C} \cup \{\infty\}$ , such that  $\pi(c) = c$ , for every  $c \in \mathbb{C}$  and  $\pi(x) = \infty$ , for every  $x \in S \setminus \mathbb{C}$ , is useful.

Another metrizable compactification is the one defined in [8] and constructed as follows: consider the map

$$\begin{aligned} \phi : \mathbb{C} &\longrightarrow D = \{\lambda \in \mathbb{C} : |\lambda| < 1\} \\ z &\longmapsto \frac{z}{1 + |z|} \end{aligned} ,$$

which is a homeomorphism. A compactification of the image  $D$  of  $\phi$  is  $\overline{D}$ , the closure of  $D$ , with the usual metric. This leads to the following compactifica-

tion of  $\mathbb{C}$

$$(2.1) \quad S_1 := \mathbb{C} \cup \{\infty e^{i\vartheta} : 0 \leq \vartheta \leq 2\pi\},$$

with metric  $d$  given by

$$(2.2) \quad \begin{aligned} d(z, w) &= \left| \frac{z}{1+|z|} - \frac{w}{1+|w|} \right| && \text{if } z, w \in \mathbb{C}, \\ d(z, \infty e^{i\vartheta}) &= \left| \frac{z}{1+|z|} - e^{i\vartheta} \right| && \text{if } z \in \mathbb{C}, \vartheta \in \mathbb{R}, \\ d(\infty e^{i\vartheta}, \infty e^{i\varphi}) &= \left| e^{i\vartheta} - e^{i\varphi} \right| && \text{if } \vartheta, \varphi \in \mathbb{R}. \end{aligned}$$

In what follows, with a *compactification*  $(S, \varrho)$  of  $\mathbb{C}$ , we shall always mean a *metrizable compactification*.

An important question for a given compactification of  $\mathbb{C}$  is, whether for  $c \in \mathbb{C}$  and  $x \in S \setminus \mathbb{C}$ , the addition  $c + x$  is well defined. In other words, having two convergent sequences  $\{z_n\}, \{w_n\}$  in  $\mathbb{C}$ , such that  $z_n \rightarrow c$  and  $w_n \rightarrow x$  does the sequence  $\{z_n + w_n\}$  have a limit in  $S$ ?

If the answer is positive for any such sequences  $\{z_n\}, \{w_n\}$  in  $\mathbb{C}$ , then the limit  $y \in S$  of the sequence  $\{z_n + w_n\}$  is uniquely determined and we write  $c + x = y = x + c$ . We are interested in compactifications  $(S, \varrho)$ , where  $c + x$  is well defined for any  $c \in \mathbb{C}$  and  $x \in S$  (it suffices to take  $x \in S \setminus \mathbb{C}$ ). In this case, the map  $\mathbb{C} \times S \rightarrow S$ ,  $(c, x) \mapsto c + x$ , is automatically continuous.

Indeed, let  $x \in S \setminus \mathbb{C}$ ,  $y \in \mathbb{C}$  and  $w = x + y \in S \setminus \mathbb{C}$ . Let  $\{z_n\}$  in  $S$  and  $\{y_n\}$  in  $\mathbb{C}$ , such that  $z_n \rightarrow x$  and  $y_n \rightarrow y$ . If all but finitely many  $z_n$  belong to  $\mathbb{C}$ , then by our assumption  $z_n + y_n \rightarrow x + y$ . Suppose that infinitely many  $z_n$  belong to  $S \setminus \mathbb{C}$ . Without loss of generality we may assume that all  $z_n$  belong to  $S \setminus \mathbb{C}$  and by compactness we can assume that  $z_n + y_n \rightarrow l \neq w = x + y$ .

Let  $d = \varrho(l, w) > 0$ . Then there exists  $n_0 \in \mathbb{N}$ , such that

$$\varrho(z_n + y_n, l) < \frac{d}{2} \quad \text{for all } n \geq n_0.$$

Fix  $n \geq n_0$ . Since,  $z_n + y_n$  is well defined, there exists  $z'_n \in \mathbb{C}$ , such that

$$\varrho(z_n, z'_n) < \frac{1}{n} \quad \text{and} \quad \varrho(z_n + y_n, z'_n + y_n) < \frac{1}{n}.$$

It follows that

$$\varrho(z'_n, x) \leq \varrho(z'_n, z_n) + \varrho(z_n, x) < \frac{1}{n} + \varrho(z_n, x) \rightarrow 0.$$

Hence,  $z'_n \rightarrow x$ ,  $y_n \rightarrow y$  and  $z'_n, y_n \in \mathbb{C}$ . By our assumption, it follows that  $z'_n + y_n \rightarrow x + y = w$ . But

$$\begin{aligned} \varrho(z'_n + y_n, l) &\leq \varrho(z'_n + y_n, z_n + y_n) + \varrho(z_n + y_n, l) \\ &\leq \frac{1}{n} + \varrho(z_n + y_n, l) < \frac{1}{n} + \frac{d}{2} \rightarrow \frac{d}{2}. \end{aligned}$$

Thus, for all  $n$  large enough we have

$$\varrho(z'_n + y_n, l) \leq \frac{3d}{4} < d = \varrho(l, w).$$

It follows that  $\varrho(z'_n + y_n, w) \geq \frac{d}{4}$ , for all  $n$  large enough. Therefore, we cannot have  $z'_n + y_n \rightarrow w$ .

Consequently, one concludes that the addition map is continuous at every  $(x, y)$  with  $x \in S \setminus \mathbb{C}$  and  $y \in \mathbb{C}$ . Obviously, it is also continuous at every  $(x, y)$  with  $x$  and  $y$  in  $\mathbb{C}$ . Thus, addition is continuous on  $S \times \mathbb{C}$ . Furthermore, the following holds:

Let  $K \subset \mathbb{C}$  be compact. Obviously, the map  $K \times S \rightarrow S$ ,  $(c, x) \mapsto c + x$ , is uniformly continuous.

*Remark 1.* The preceding certainly holds for the compactification  $(S_1, d)$  (see (2.1)), since

$$c + \infty e^{i\vartheta} = \infty e^{i\vartheta} \quad \text{for all } c \in \mathbb{C} \text{ and } \vartheta \in \mathbb{R},$$

and we have continuity.

*Remark 2.* If we identify  $\mathbb{R}$  with the interval  $(-1, 1)$ , up to a homeomorphism, then  $\mathbb{C} \cong \mathbb{R}^2$  is identified with the square  $(-1, 1) \times (-1, 1)$ . An obvious compactification of  $\mathbb{C}$  is then the closed square with the usual metric. The points at infinity are those on the boundary of the square, for instance, those points on the side  $\{1\} \times [-1, 1]$ . If  $x \in \{1\} \times (-1, 1)$  and  $c \in \mathbb{C}$ , then  $c + x$  is a point in the same side; if  $\text{Im } c \neq 0$ , then  $c + x \neq x$ . If  $x = (1, 1)$  and  $c \in \mathbb{C}$ , then  $x + c = x$ . If  $\text{Im } c > 0$ , then  $c + x$  lies higher than  $x$  in the side  $\{1\} \times (-1, 1)$ .

In this example, the addition is well defined and continuous, but the points at infinity are not stabilized as in Remark 1.

**QUESTION.** Is there a metrizable compactification of  $\mathbb{C}$  such that the addition  $c + x$  is not well defined for some  $c \in \mathbb{C}$  and  $x \in S \setminus \mathbb{C}$ ?

The answer is “yes”. An example comes from the previous square in Remark 2, if we identify all the points of  $\{1\} \times [-\frac{1}{2}, \frac{1}{2}]$  and make them just one point.

### 3. RUNGE AND MERGELYAN TYPE THEOREMS

In this section using a compactification of  $\mathbb{C}$  satisfying all properties discussed in the Preliminaries, we obtain the following theorem, that extends [5, Theorem 3.3].

**THEOREM 3.1.** *Let  $\Omega \subset \mathbb{C}$  be a bounded domain, whose boundary consists of a finite set of pairwise disjoint Jordan curves. Let  $K = \overline{\Omega}$  and  $A$  a set containing one point from each component of  $(\mathbb{C} \cup \{\infty\}) \setminus K$ . Let  $(S, \varrho)$  be a compactification of  $\mathbb{C}$ , such that the addition  $+: \mathbb{C} \times S \rightarrow S$  is well defined. Let  $f: K \rightarrow S$  be a continuous function, such that  $f(\Omega) \subset \mathbb{C}$  and  $f|_{\Omega}$  is holomorphic. Let  $\varepsilon > 0$ . Then, there exists a rational function  $R$  with poles only in  $A$  and such that  $\varrho(f(z), R(z)) < \varepsilon$ , for all  $z \in K$ .*

*Proof.* If  $\Omega$  is a disk, the proof has been given in [1]. If  $\Omega$  is the interior of a Jordan curve, the proof is given again in [1], but also in [6]. In the general case, we imitate the proof of [5, Theorem 3.3]. Namely, we consider the Laurent decomposition of  $f$ , given by  $f = f_0 + f_1 + \dots + f_N$  (see [4]). The function  $f_0$  is defined on a simply connected domain, bounded by a Jordan curve, and it can be uniformly approximated by a polynomial or a rational function  $R_0$  with pole in the unbounded component. Similarly,  $f_1$  is approximated by a rational function  $R_1$  with pole in  $A$  and so on. Thus, the function  $R_0 + R_1 + \dots + R_N$  approximates, with respect to  $\varrho$ , the function  $f = f_0 + f_1 + \dots + f_N$ . This is due to the fact that at every point  $z$  all the  $f_i$ 's,  $i = 1, 2, \dots, N$ , except maybe one, take values in  $\mathbb{C}$  and the one, maybe has as a value, an infinity point in  $S \setminus \mathbb{C}$ . In this way, the addition map  $\mathbb{C} \times S \rightarrow S$ ,  $(c, x) \mapsto c + x$ , is well defined and uniformly continuous on compact sets and so we are done. ■

Another Runge-type theorem is the following, where we do not need any assumption for the compactification  $S$ , or the addition map  $+: \mathbb{C} \times S \rightarrow S$ .

**THEOREM 3.2.** *Let  $\Omega \subset \mathbb{C}$  be open,  $f: \Omega \rightarrow \mathbb{C}$  be holomorphic and  $(S, \varrho)$  a compactification of  $\mathbb{C}$ . Let  $A$  be a set containing one point from each component of  $(\mathbb{C} \cup \infty) \setminus \Omega$ . Let  $\varepsilon > 0$  and  $L \subset \Omega$  compact. Then, there*

exists a rational function  $R$  with poles in  $A$ , such that  $\varrho(f(z), R(z)) < \varepsilon$  for all  $z \in L$ .

*Proof.* Clearly the subset  $f(L)$  of  $\mathbb{C}$  is compact. Then, from the classical theorem of Runge, there exist rational functions  $\{R_n\}$ , with poles only in  $A$ , converging uniformly to  $f$  on  $L$ , with respect to the Euclidean metric  $|\cdot|$ . Hence, there is a positive integer  $n_0$  and a compact  $K$ , such that

$$f(L) \subset K \subset \mathbb{C} \quad \text{and} \quad R_n(L) \subset K \quad \text{for all } n \geq n_0.$$

But on  $K$  the metrics  $|\cdot|$  and  $\varrho$  are uniformly equivalent. Therefore,  $R_n \rightarrow f$  uniformly on  $L$ , with respect to  $\varrho$ . To conclude the proof, it suffices to put  $R = R_n$ , for  $n$  large enough. ■

Theorem 3.2 easily yields the following

**COROLLARY 3.3.** *Under the assumptions of Theorem 3.2 there exists a sequence  $\{R_n\}$  of rational functions with poles in  $A$ , such that  $R_n \rightarrow f$ ,  $\varrho$ -uniformly, on each compact subset of  $\Omega$ .*

*Remark.* According to Corollary 3.3, some of the  $\varrho$ -uniform limits, on compacta, of rational functions with poles in  $A$ , are the holomorphic functions  $f : \Omega \rightarrow \mathbb{C}$ . Those are limits of the finite type. The other limits of sequences  $\{R_n\}$  as above may be functions  $f : \Omega \rightarrow S \setminus \mathbb{C}$  of infinite type, continuous (but maybe not all of them, as the Example  $(S_1, d)$  shows; cf. [8]).

**QUESTION.** Is a characterization possible for such limits  $f : \Omega \rightarrow S_1 \setminus \mathbb{C}$ ?

An imitation of the arguments in [8, p. 1007] gives that  $f$  must be of the form  $f(z) = \infty e^{i\vartheta(z)}$ ,  $z \in \Omega$ , where  $\vartheta$  is a multivalued harmonic function.

The following extends [5, Section 5].

**THEOREM 3.4.** *Let  $\Omega \subset \mathbb{C}$  be open and  $f$  a meromorphic function on  $\Omega$ . Let  $B$  denote the set of poles of  $f$ . Let  $(S, \varrho)$  be a compactification of  $\mathbb{C}$ , such that the addition  $+: \mathbb{C} \times S \rightarrow S$  is well defined. Let  $\varepsilon > 0$  and  $K \subset \Omega$  be a compact set. Then, there is a rational function  $g$ , such that  $\varrho(f(z), g(z)) < \varepsilon$ , for every  $z \in K \setminus B$ .*

*Proof.* Since  $B \cap K$  is a finite set, the function  $f$  decomposes to  $f = h + w$ , where  $h$  is a rational function with poles in  $B \cap K$  and  $w$  is holomorphic on an open set containing  $K$ . By Runge's theorem there exists a rational function  $R$

with poles off  $K$ , such that  $|w(z) - R(z)| < \varepsilon'$  on  $K$ . Since  $w(K)$  is a compact subset of  $\mathbb{C}$  and the addition  $+: \mathbb{C} \times S \rightarrow S$  is well defined, a suitable choice of  $\varepsilon'$  gives

$$\varrho([h(z) + w(z)], [h(z) + R(z)]) < \varepsilon \quad \text{on } K \setminus B.$$

We set  $g = h + R$  and the result follows. ■

#### 4. ARAKELIAN SETS

A closed set  $F \subset \mathbb{C}$  is said a *set of approximation* if every function  $f : F \rightarrow \mathbb{C}$  continuous on  $F$  and holomorphic in  $F^\circ$  can be approximated by entire functions, uniformly on the whole  $F$ . This is equivalent to the fact that  $F$  is an Arakelian set (see [2]), that is  $(\mathbb{C} \cup \{\infty\}) \setminus F$  is connected and locally connected (at  $\infty$ ).

We can now ask about an extension of the Arakelian theorem in the context of metrizable compactifications. A result in this direction is the following

**PROPOSITION 4.1.** *Let  $F \subset \mathbb{C}$  be a closed Arakelian set with empty interior, i.e.,  $F^\circ = \emptyset$ . We consider the compactification  $(S_1, d)$  of  $\mathbb{C}$  (see (2.1) and (2.2)) and let  $f : F \rightarrow S_1$  be a continuous function. Let  $\varepsilon > 0$ . Then, there is an entire function  $g$  such that  $d(f(z), g(z)) < \varepsilon$ , for every  $z \in F$ .*

*Proof.* According to (1.1), the compactification  $S_1$  is homeomorphic to  $\overline{D} = \{z \in \mathbb{C} : |z| \leq 1\}$ . For each  $0 < R < 1$  let us define

$$\begin{aligned} \phi_R : \overline{D} &\longrightarrow \{z \in \mathbb{C} : |z| \leq R\} \subset \overline{D} \\ z &\longmapsto \begin{cases} z, & \text{if } |z| \leq R, \\ \frac{Rz}{|z|}, & \text{if } R \leq |z| \leq 1. \end{cases} \end{aligned}$$

In other words, the whole line segment  $[Re^{i\vartheta}, e^{i\vartheta}]$  is mapped at the end point  $Re^{i\vartheta}$ . The function  $\phi_R$  is continuous and induces a continuous function  $\tilde{\phi}_R : S_1 \rightarrow S_1$ . It suffices to take  $\tilde{\phi}_R := T^{-1} \circ \phi_R \circ T$ , where  $T : S_1 \rightarrow \{w \in \mathbb{C} : |w| \leq 1\}$  is defined as follows

$$\begin{aligned} T(z) &:= \frac{z}{1 + |z|} && \text{for } z \in \mathbb{C} \subset S_1, \\ T(\infty e^{i\vartheta}) &:= e^{i\vartheta} && \text{for } \vartheta \in \mathbb{R}. \end{aligned}$$

If  $\varepsilon > 0$  is given, then there exists  $R_\varepsilon < 1$ , such that for  $R_\varepsilon \leq R < 1$  and  $z \in S_1$ , we have  $d(z, \tilde{\phi}_R(z)) < \frac{\varepsilon}{2}$ .

Let now  $f$  be as in the statement of the Proposition 4.1. Then,

$$d\left(f(z), (\tilde{\phi}_R \circ f)(z)\right) < \frac{\varepsilon}{2} \quad \text{for all } z \in F.$$

Moreover, the function  $\tilde{\phi}_R \circ f : F \rightarrow \mathbb{C}$  is continuous. Since  $F$  is a closed Arakelian set, with empty interior, and  $(\tilde{\phi}_R \circ f)(F) \subset K$ , is included in a compact subset  $K$  of  $\mathbb{C}$ , there exists  $g$  entire, such that

$$\left|(\tilde{\phi}_R \circ f)(z) - g(z)\right| < \varepsilon' \quad \text{for all } z \in F.$$

Since  $(\tilde{\phi}_R \circ f)(F)$  is contained in a compact subset  $K$  of  $\mathbb{C}$ , for a suitable choice of  $\varepsilon'$ , it follows that

$$d\left((\tilde{\phi}_R \circ f)(z), g(z)\right) < \frac{\varepsilon}{2} \quad \text{for all } z \in F.$$

The triangle inequality completes the proof. ■

An analogue of Proposition 4.1 for the one point compactification  $\mathbb{C} \cup \{\infty\}$  of  $\mathbb{C}$  has been established in [5].

#### REFERENCES

- [1] I. ANDROULIDAKIS, V. NESTORIDIS, Extension of the disc algebra and of Mergelyan's theorem, *C.R. Math. Acad. Sci. Paris* **349**(13–14) (2011), 745–748.
- [2] N.U. ARAKELIAN, Uniform approximation on closed sets by entire functions, *Izv. Akad. Nauk SSSR Ser. Mat.* **28** (1964), 1187–1206 (Russian).
- [3] L. BROWN, P.M. GAUTHIER, W. HENGARTNER, Continuous boundary behaviour for functions defined in the open unit disc, *Nagoya Math. J.* **57** (1975), 49–58.
- [4] G. COSTAKIS, V. NESTORIDIS, I. PAPADOPERAKIS, Universal Laurent series, *Proc. Edinb. Math. Soc. (2)* **48** (3) (2005), 571–583.
- [5] M. FRAGOULOPOULOU, V. NESTORIDIS, I. PAPADOPERAKIS, Some results on spherical approximation, *Bull. Lond. Math. Soc.* **45** (6) (2013), 1171–1180.
- [6] V. NESTORIDIS, Compactifications of the plane and extensions of the disc algebra, in “Complex Analysis and Potential Theory”, CRM Proc. Lecture Notes, 55, Amer. Math. Soc., Providence, RI, 2012, 61–75.
- [7] V. NESTORIDIS, An extension of the disc algebra, I, *Bull. Lond. Math. Soc.* **44** (4) (2012), 775–788.

- [8] V. NESTORIDIS, N. PAPADATOS, An extension of the disc algebra, II, *Complex Var. Elliptic Equ.* **59** (7) (2014), 1003–1015.
- [9] V. NESTORIDIS, I. PAPADOPERAKIS, A remark on two extensions of the disc algebra and Mergelian's theorem, *preprint* 2011, arxiv: 1104.0833.
- [10] W. RUDIN, "Real and Complex Analysis", McGraw-Hill Book Co., New York-Toronto, Ont.-London, 1966.